UNCLASSIFIED

AD NUMBER AD840889 NEW LIMITATION CHANGE TO Approved for public release, distribution unlimited **FROM** Distribution authorized to U.S. Gov't. agencies and their contractors; Foreign Government Information; 12 JUL 1967. Other requests shall be referred to US Army Biological Laboratory, Attn: Technical Release Brancn [TID], Frederick, MD 21701. **AUTHORITY** SMUFD d/a ltr, 15 Feb 1972

TRANSLATION NO. 1980

DATE: 12 Yuly 1967

DDC AVAILABILITY NOTICE

Reproduction of this publication in whole or in part is prohibited. However, DDC is authorized to reproduce the publication for United States Government purposes.

STATEMENT #2 UNCLASSIFIED This document is subject to special export controls and each transmittal to foreign controls and each transmittal to roeign made governments or foreign nationals may be made governments or foreign nationals may be made only with prior approval of Dept. Of Army, Technical Release Branch Fort Detrick, ATTN: Technical Release Branch TID Erederick Maryland 21701 TID. Frederick, Maryland 21701



DEPARTMENT OF THE ARMY Fort Detrick Frederick, Maryland

BRONCHOSCOPICAL OBSERVATIONS AFTER INHALATION OF DES UNDER NORMAL AND MORSID CONDITIONS OF THE RESPIRATORY TRACT

p. 2-3

Dr W. Schiessle, Freiburg im Breisgau, Medical University Clinic, Internal Department of the Robert Koch Clinic (Director: Professor Dr. and Dr. hon. causa L. Heilmeyer)

The introduction gives a brief survey of the methods hitherto used in order to investigate the penetration of the respiratory tract of enimals and man by inhaled aerosols.

A report is then given on the methodology of the author's own dye-inhalation experiments on man. A small number of persons in the experiment inhaled Evans Blue with a bronchoscope inserted in the upper trachea or in the lower interbronchial area. Most of them inhaled the dye before bronchoscopy.

In demonstrating the results in the normal respiratory tract and in case of various pathological conditions in the bronchial system and in the pulmonary parenchyma, the author in each instance shows first of all a photograph giving a general view of the lungs, often supplemented by photographs of the layers, and then the bronchoscopically obtained color photograph with the dye precipitation that is present, usually supplemented by bronchograms.

In the discussion of the results the conditions for the precipitation of inhaled dye and the intensity thereof are discussed on the basis of schematic diagrams. Conclusions as to the possibilities and limits of aerosol therapy are drawn. Also dealt with it the significance of bronchoscopic investigations for clustifying further scientific questions of serosol inhalation.